

Frequently Used Compounds: Gastroenterologist

- **Promethazine 12.5 mg/0.1mL Gel**
Sig: Apply 0.1 – 0.2 mL to inner wrist every 4 – 6 hours as needed
- **Diltiazem 2% Cream**
Sig: Apply pea-sized amount to peri-anal area twice daily
- **Nitroglycerin 0.2% Ointment**
Sig: Apply pea-sized amount to peri-anal area twice daily
- **Nifedipine 0.2% Ointment**
Sig: Apply pea-sized amount to peri-anal area twice daily
- **Rocket Lidocaine 2% – Hydrocortisone 1% suppository**
Sig: one suppository at bedtime
- **Phenylephrine 30% Ointment or 40% Ointment**
Sig: Apply pea-sized amount to peri-anal area once to twice daily
- **Vancomycin Liquid (Oral) 250mg/5mL**
Sig: _____
- **Metronidazole 10% Ointment**
Sig: _____
- **Naltrexone 4.5mg Capsules (for Crohn’s Disease)**
Sig: 1 capsule at bedtime
- **Naltrexone 0.5mg Capsules (for IBS)**
Sig: 1 capsule at bedtime
- **Ketoprofen 10% Lipoderm Gel (Other strengths available, please call)**
Sig: Apply to affected area 2 – 4 times daily



Formulas may be changed by the Prescriber in any way as to meet the specific needs of the individual!

T&C-FUC-Gastro-ChrisC-revJH-0512-0115

106 Prospect Street, Ridgewood, NJ 07450 Ofc: 201-447-2020 Fax: 201-447-3253

www.tccompound.com

info@tccompound.com