

Vasceptor Sublingual/Topical Gel Order Form

Patient Name		Prescriber Name	
Date of Birth		Office Phone #	
Phone #		Office Fax #	
Street Address		Street Address	
City, State, Zip		City, State, Zip	
Allergies		NPI	
Email		DEA	
<input type="checkbox"/> Send to PATIENT <input type="checkbox"/> Charge PATIENT		<input type="checkbox"/> Send to OFFICE <input type="checkbox"/> Charge OFFICE	

Vasceptor® Curcuminoid Sublingual/Topical Gel, 8.5%

Preferred Sublingual Directions (Please check off all that applies):

☐ **Sublingual:** Apply 1-2 clicks (0.05-0.1mL) on spoon or finger with or without a glove. Vigorously rub it under the tongue for 15-30 seconds. Wait 1-2 minutes before spitting it out, swallowing or drinking. Wash the stained finger with a high fat (lanoline) containing soap such as Dove. This procedure can be repeated twice daily if needed.

☐ **Alternative Topical Directions (Please check off all that applies):**

☐ **Knee/Shoulder:** Apply 2 clicks (0.1mL) and rub thoroughly into the soft parts below and on the sides of the kneecap. Use once daily as long as needed.

☐ **Hand/Wrist/Elbow/Muscle Strain:** Apply 1-2 clicks (0.05-0.1mL) at the site of pain and rub in thoroughly. Use once daily as long as needed.

☐ **Back:** Apply 2-4 clicks (0.1-0.2mL) at the site of pain and rub in thoroughly. Use once daily as long as needed.

Quantity:

☐ 1 mL (1 unit) \$35

☐ 5mL (1 unit) \$150

☐ 9mL (1 unit) \$240

Refills: __ ☐ 1 year

Prescriber Signature: _____

Date Written: __ / __ / ____

Toll-Free Phone: 800-850-2101

Phone: 201-447-2020

Fax: 201-447-3253

Pharmacist@tccompound.com

www.tccompound.com

535 East Crescent Avenue Ramsey, NJ 07446